Willey: Outline of Taylor's Flora Antarctica Vol. 1

Seems to be the handwriting
of Henry Willey (1824-1907)
of New Bedford, Mass.

Flora Antarctica Vol. I. 1844

(Auckland & Campbell's Islands)

Lichenes

(By Dr Thos Taylor & J. D. Hooker)

Usnea plicata Ach.

" hirta "

" barbata Ach.

" Sulphurea "

Ramalina inflata, Hook. fil. et Taylor.

(Cetraria inflata in Lon. Jour. Bot.

Sphaerophoron tenerum Laur.

" " " coralloides (S. coralloides in Lon Journ. of Bot.)

" australe Laur.

" compressum Ach.

Stereocaulon paschale Ach.

" Hook. fil. et Tayl.

2.

Ösmonyée Bonifácius Sek.

" Aggregata " (Infusoria stolata -
- Lor. Jam Bol.)

" Semocysta Margaritá Sek.
(L. samentöte i Lor. Jam Bol.)

" Hyphidea Sek.

" " c. rigida (L. rigida i L. J. B.)

" Mucedó Sek.

Pstricha polygastrica Sek.

Stricta oligomera Sek.

" Familia Oxlicae (S. linearis typus i
L. J. B.)

" Hygienetii Oxlicae. (S. flabra i L. J. B.)

" Cellulifera Stark. fälet-Zayl.

" Menzáesii " " " "

" Richardii Mont.

Parmelia subsinuosa Sch.

" Sphinctrina turbinata (" Lindly distinct from P. subsinuosa ").

Lecanora hyporeum Sch.

" virescens. Hook. fil & Tayl. (affinis L. varia)

" testacea. Sch.

" Pontica Sch.

Lecidea geomaea Tayl. in Hook. Icon.

" incana. Hook. (Beltona parhygrophila

Porina granulata Hook. fil & Tayl.

Thelotrema leptoleucum Sch.

Opegrapha alba Pers.

(Fuegia. Ile Falklands. etc.)

Lichens

Usnea melaxantha Ach.

" v. a Beltonii;

v. β. Jacquinii;

v. γ. fasciata (U. fasciata Lowe)

v. δ. sphaculata (U. sphaculata Br.)

Hab. Throughout these Isles from the sea to

Lowest mast-attainment

" Lyellii. Hook. fil. (U. melaxantha similima)

" barbata. Ach.

" v. articulata Ach.

" plicata Ach.

" floridea Ach.

Evernia Magellanica. Mont.

Ramalina scopulorum Ach. cum var.

Cetraria Islandica Ach.

" glauca Ach.

6.

Asteria sepenicola Ach.

" accolata Fr.

Bacidia polorina Ach.

" cellulosa "

Belonia polydactyla Ach.

" conica " ?

" venosa " ?

" horizontalis "

Biatora crocea L.

" " v. Bjelkeana (S. imperf. in L. L. B.)

" endochrysa, Delise.

" oxygmaea. Ach.

" obvoluta "

" Billotiana, Delise = B. Richardi Fr. (Jaccard Fl.

" " v. B. (Schleicheri Fr. in L. C.

" fauvolata Delise.

" vernalis. Ach.

Stella Thomasii Delise. (S. senticulata in L. f. B.

" Bryoniae " (S. globus in L. f. B.)

" Jelinii Bek.

Stenocaulon corallinum Fr. (S. pascale in L. f. B.)

" ramulosum Bek.

" alpinum Fr.

" dendroideum Somm.

Stereocaulon coralloides Bek.

" tenuem, Lauer.

" compressum, Bek.

" austrule, Laur.

" frozile, Bek.

Cenomyce pyxidata Bek.

" gracilis "

" fimbriata "

" v. subulata (C. subulata in L. f. B.

" pertusata Bek.

" cornuta "

" " v. Thomasii Delise

8

Cenomyce furcata Sek.

" " v. squamulosa Sek.

" coccifera Sek.

" deformis Sek.

" sorediosa "

cum vers. strepsilis & sylvatica

" uncialis Sek.

" ? verniculoris, Sek.

" aggregata Sek.

" bacillaris "

" sparassa Sek.

Parmelia enteromorpha. Sek.

" " v. dendrita

" dendrytes Sek.

" encausta "

" saxatilis "

" subaurifera "

" v. sphenetrina

(P. " Mark.)

Leucoium candidum Bert.?

" vernum " (abundant=)

" autumnale "

" bulbocastanum ?

" Spilotii Fr.

" contiguum Fr.

" alno-album Bert.

" turgidum Lonnf.

" fuscatum Bert.

" stellulatum Foyl.

" floccosum Schr.

" geographicum "

" pondosum Bert.

" tabulatorum "

" comidops Fr. ?

" orbicum Lonnf.

" miliare v. leprosum Fr.

" obscurum - in rubens Fries.

Gyrophora cruca v. Schr. (non v. hirsuta)

12.

Hypnopsis alata Pers.

Sirtonia polymorpha Sch.

Parmelia communis DC. abundant.

" Wulfenii DC.

Phlyctina hyperbolina Sch.

Verrucaria umbrina Sch. (v. selida in L. f.
L. J. B.)

Collema crispum Sch. (dispersa)

" hemitrichum " ?

" Solenianum "

" v. anastale (Caustale in L.
in L. J. B.